
NUTRITIONAL POTENTIALS AND CHEMICAL VALUE OF SOME TROPICAL LEAFY VEGETABLES CONSUMED IN SOUTH WEST NIGERIA

Sobowale, S.S^{1*}; Olatidoye, O.P²; Olorode, O.O¹, and Akinlotan, J.V²
¹*Department of Food Science and Technology, Moshood Abiola polytechnic, Nigeria*
²*Department of Food Science and Technology, Igbinedion University Okada, Nigeria*
Email: sunsam_2003@yahoo.co.uk

ABSTRACT

The human population in tropical African depends largely on a number of edible leafy vegetables to meet up with shortages in minerals and vitamins. Data base on the nutritional value of these leafy vegetables is scanty. Therefore, we report the nutritional components in three species of leafy vegetables using standard analytical methods. All the vegetables contained moisture (79.92 to 84.0%), crude protein (20.61 to 22.7%), crude fibre (10.7 to 22.44%), ash (6.8 to 10.44%), carbohydrate (55.86 to 68.22%) crude lipid (4.24 to 5.6%) and food energy (1507.19 to 1673.96 kJ/100g). The mineral element content were high with remarkable concentration of K(35.2 to 48.8mg/100g),Na(11.4 to 14.4 mg/100g),Ca(15.4 to 18.7mg/100g), Mg(12.2 to 18.7mg/100g), P(13.8 to 15.08mg/100g). The relationship between Na and K as well as between Ca and P; are desirable with the respective ratios of Na/K (0.6) and Ca/P (1.2). They also contain high levels of carotenoids (30 to 41.5 mg/100g DW), vitamin C (137.5 to 197.5 mg/100g DW). Seventeen amino acids (isoleucine, leucine, lysine, methionine, cysteine, phenylalmine, tyrosine, threonine, valine, alanine, arginine, aspartic acid, glutamic acid, glycine, histidine, proline and serine) were detected. Their amino acid composition compare favourably with that of WHO/FAO protein standard indicating favourable nutritional balance except for lysine and methionine which appear marginal. The nutritional values of the phytochemicals were also assessed with a view of establishing and understanding their nutritional uses. The functional properties for the three vegetables were similar. Comparing the nutrient and chemical constituents with recommended dietary allowance (RDA) values, the results reveal that the leaves contain an appreciable amount of nutrients, minerals, vitamins, amino acids and phytochemicals and low levels of toxicants.

Keywords: vegetables, nutrients, amino-acids, phytochemicals, functional properties

INTRODUCTION

The most serious threat to survival of humanity is the increasing gap between population growth and food supply. It has been estimated that over 500 million people in the world today are malnourished [1]. Unfortunately an estimated 789 million people in developing countries still suffer from malnutrition especially infants and children of rural areas [2]. In Nigeria, as in most other tropical countries of Africa where the daily diet is dominated by starchy staple foods, vegetables are the cheapest and most readily available sources of important proteins, vitamins, minerals and essential amino acid [3]. Green leafy vegetables are succulent plant grown in the gardens and consume as side dish or soup with foods among the Nigerian tribes. Vegetables have been identified to contain high content of mineral, antioxidant and vitamins [4]. Organoleptically, vegetables are valued for their

supreme flavor and aroma, crop texture, attractive colour and their overall appeal to human sense of smell, taste, touch, and sight [5]. They contribute considerable amount of vitamins A, C, B₆, thiamin, niacin, riboflavin and minerals which are essential to metabolic process in the body system in addition they supply proteins, starch and sugars and are important source of dietary and crude fibre. [6, 7, 8, 9, 10, 11, 12]. Many of the local vegetable materials are under-exploited because of inadequate scientific knowledge of their nutritional potentials. The lack of nutritional information and inadequate development of nutritionally improved products from local raw materials have direct bearing on nutrition. In order to arrest the situation, much attention has been centered on the exploitation and utilization of non-conventional plant material for food [13].

Several species of these vegetables abound in Nigeria and many West African countries where they are used as condiments or spices in human diets, condiments or supplementary feeds to livestock such as rabbit, swine and cattle [14, 15]. Though several works reporting compositional evaluation and functional properties of various types of edible plants in use in the developing countries abound in literature, much still need to be done. Many workers [16, 17, 18, 19, 20, 21, 22, 23, 24] have reported the compositional evaluation of various types of leafy vegetables in use in the developing countries. Several of these vegetables species abound in Nigeria, *O. gratissimum*, *T. occidentalis*, *V. amygdalina* are known to be amongst the leafy vegetables consumed in Nigeria. Their nutritional contribution has not been widely exploited hence nutritional information on these species of vegetables will be useful for the nutritional education of the public and to improve the nutritional status of the population. As a result of the beneficial attributes of these leafy vegetables, there is need to establish chemical, nutritional and toxicological properties before advocating for increased utilization.

MATERIALS AND METHOD

Preparation of samples: Fresh leaves of some of the vegetables species were harvested from an experimental farm in Abeokuta. They were rinsed in water to remove dust, sorted and sliced. Some of the fresh samples were used for moisture determination while the rest of the samples were dried at 45⁰C to constant weight using a Gallenkamp oven. They were ground to fine power in a sieve through 1.0mm mesh and stored in airtight container for laboratory analysis.

Chemical analysis: Proximate composition, dietary fibre, total carotenoids and gross energy value of the vegetable leaves were determined by method of the Association of official chemists [25] while mineral like sodium and potassium content were determined by flame photometry, phosphorus were determined by the vonodomobybdate method [25]. Other mineral were determined after wet digestion using Atomic Absorption spectrophotometer (AAS model, SP9). Amino acid analysis was carried out using amino acid analyzer.

Determination of antinutrient: Oxalic acid, Saponin. Tannin content of the raw and processed vegetables samples were determined according methods described by AOAC [25].

Statistical analysis: All data obtained in triplicate were subjected to analysis of variance (ANOVA) using the SPSS statistical package (version 10.0), (2000 edition).

RESULTS AND DISCUSSION

The proximate composition of the three leafy vegetables is shown in Table1 with *O. gratissimum* having the highest moisture content of 84.00% followed by *V.amygdalin* with a value of 82.12% and *T. occidentalis* with 80.88%. The high level of moisture in all the samples investigated suggests that the leafy vegetables would not store for long without spoilage since a higher water activity could enhance microbial action bringing about food spoilage. It has been reported that fruits and vegetables contain as high as 85% water [26] consistent with the range of moisture level obtained in this study. The crude protein range of 5.7-21.02% for the leaves investigated and was relatively higher than some foodstuffs like *L. sativa* (0.7-1.1%), *Portulaca* leaves (1.3-1.5%), cassava leaves (7.1-8.2%) [27]. Values for these samples are close to values obtained by Igile [28] for *V. amygdalina*, Ejoh [11] for *V.amygdalin* and Fevrier and Viroben [29] for other leafy vegetables. The potentials of these vegetables as food or alternative protein source are shown by their protein and mineral composition. Aletor and Adeogun [15] opined that the use of these vegetables in amino acid supplementation in meeting protein and energy requirement could be a beneficial factor. The supplementation of plant foods with other food can give rise to high biological value and satisfactorily meeting the protein needs of adults [29]. The protein in leafy vegetables would require dietary supplementation with cereal protein and legume to alleviate kwashiorkor. The crude fibre was highest in *V.amygdalin* with 22.44g/100gdw and lowest in *O. gratissimum* at 10.7g/100gdw. These vegetables have high level of dietary fibre which varied significantly in the different species. Craplet and craplet-Meuner [30] and Tanya et al., [31] affirmed that leafy vegetables are rich in dietary fibre. Eun-Hee et al., [32] opined that the average level of dietary fibre in leafy vegetables in Asian countries to be 33% dry weight. This result is in close agreement with work [11, 32].

High level of dietary fibre in leafy vegetables are advantageous for their active role in the regulation of intestinal transit, increasing dietary bulk and increasing faeces consistency due to their ability to absorb water[33,34]. Ash content was relatively high with values ranging from 10.44% in *T. occidentalis* to 6.8% in *O. gratissimum*. These values except for *O.gratissimum* were found to be higher than values obtained for *Amaranthus* species [35]. These values obtained is higher than the values obtained by [11] for *V.amygdalin* and in close agreement with the work of Fasuyi [36] for *Telfairia occidentalis*. These values are an indication that these vegetables species are good sources of minerals when compared to values obtained for cereals and tubers [37]. Ether extract (fat) had a wide range of 7.62% in *Telfairia occidentalis* to 1.08% in *O.gratissimum*. Igile [28] and Ejoh [11] reported a similar value for *V. amygdalina* while Fasuyi [36] reported similar values for *T. occidentalis*. Singhai and Kulkarni [35] reported higher values for *Amaranthus* spp. These values for fat in these vegetables species confirmed the findings of many authors which showed that leafy vegetables are poor sources of lipids [11]. However, their consumption in large amount is a good dietary habit and may be recommended to individuals suffering from obesity.

Carbohydrate content of these leafy vegetables was relatively high with *T. occidentalis* having 55.86% and *V. amygdalina* having 68.22%. These carbohydrate sources are not used in the body because most of them remain undigested. Igile [28] obtained slightly higher value for *v. amygdalina* while Ejoh [11] obtained slightly lower values. The differences may be due to the physiological state of the plant before harvesting, climate and agricultural practices [35].

The gross energy values of these vegetables species are in the range of 392.67 in *V. amygdalina* to 44.8 in *ocimum gratissimum*. These levels are low due to low crude fat and relatively high level of moisture. The daily energy requirement of 2500 to 3000 kCal has been reported for adult [37]. Hence, individual would need to consume 700g of *V. amygdalin* to obtain an energy value of 2750 kcal per day which is within the range reported by Bingham [38]. When one considers the roles of protein, fiber, Ca, Fe, total carotenoids and ascorbic acid in human nutrition, the average nutritive value of the *T. occidentalis* is superior to those the other species (Table 3). These values are lower than values obtained by Kuti and Kuti [39] for different species of spinach. These results are close to those reported by Ejoh [11] for other leafy vegetables.

Table 4 shows the amino acid analysis of the three leafy vegetables (g/100 g protein) .The major abundant amino acids were Glutamic acid, Arginine, Aspartic acid and Leucine with the values of 15.2, 5.8, 5.2 and 5.1 g/100 g protein respectively. This observation is in close agreement with the work of [40, 41, 42, 43]. The sum of the aspartic acid and glutamic amino acid was 21.2 g/100 g protein. This value is lower than values obtained from selected oil seeds (melon, pumpkin, cashew nut flour and gourd seeds) ranging between 24.2-29.5 [40, 43, 44.]. The Total Amino Acid (TAA) of 34.8 g/100 g protein indicated that these vegetables will contribute significantly to the supply of amino acid in diet. This value is lower than that of melon, pumpkin and gourd seed of 53.4, 38.3 and 53.6 g/100 g protein respectively reported by [40]; soybean, 44.4 g/100 g protein [44]; Pigeon peas, 45.2 g/100 g protein [23]. This is an indication that vegetables are good sources of essential amino acids. The amino acid requirements mg/kg/day for infants and 10-12 year old children have been reported to be 31 for Isoleucine, 64 for lysine, 27 for sulfur amino acids, 37 for threonine and 14 for tryptophan [45]. There are also evidences that histidine is an essential amino acid for adults and its dietary requirement in both normal and uremic men has been reported to be between 8 and 12 mg/kg/day [46, 47]. However, Wallece [48] found that the essential amino acids were lower than the FAO/ WHO reference protein in some non conventional leafy vegetables. Comparatively most of the amino acid profile was similar for all the different species of leafy vegetables. However *V. amygdalina* had slightly lower essential amino acids is evident as values of isoleucine, leucine and valine were lower for the other samples. Histidine, tyrosine and proline were highest in *T. occidentalis* while *V. amygdalina* has the highest level of aspartic acid, glycine and methionine. Deosthale [49] showed that excess leucine in foods interfered with the utilization of isoleucine and lysine.

For all the species, the levels are generally high and stand the risk of complexing with minerals such as calcium [50]. Oke [51] in Nigeria obtained similar observations. Similar

values in amaranth [52] and spinach leaves [53] have been reported earlier. [54] reported a toxic level for oxalic acid at 3g per day. Polyphenol values ranged from 5.52mg/100g dry weight for *T. occidentalis* to 10.12mg/100g dry weight for *O. gratissimum*. Higher polyphenols content in fenugreek leaves and lower in amaranth leaves have been reported earlier [Gupta *et al.*, 1989; 55]. Saponin values for these species ranged from 0.57 in *O. gratissimum* to 1.94mg/100g dry weight in *V. amygdalina*. Saponin in foods can be astringent [56] or not bitter [57]. Saponins in plants have been known to protect the plant from fungal and insect attacks [56]. Other studies have proven that saponin have the ability to reduce the cholesterol levels in man and animals [57]. According to Ladeji [58], oxalate can bind to calcium present in food thereby rendering calcium unavailable for normal physiological and biochemical role such as the maintenance of strong bone, teeth, cofactor in enzymatic reaction, nerve impulse transmission and as clotting factor in the blood. The calcium oxalate, which is insoluble, may also precipitate around soft tissues such as the kidney, causing kidney stones [Oke, 1969]. Though loss of calcium leads to degeneration of bones, teeth and impairment of blood clotting process [59], the values obtained for these fruits were below the established toxic level. According to Oke [1966], a phytate diet of 1-6% over a long period decreases the bioavailability of mineral elements in mono gastric animals. Phytic acid can bind to mineral elements such as calcium, zinc, manganese, iron and magnesium to form complexes that are indigestible, thereby decreasing the bioavailability of these elements for absorption [60]. Phytic acid also has a negative effect on amino acid digestibility thereby posing problems to non-ruminant animals due to insufficient amount of intrinsic factor phytase necessary to hydrolyze the phytic acid complexes [61]. Phytate is also associated with nutritional diseases such as rickets and osteomalacia in children and adults respectively. High saponin level has been associated with gastroenteritis manifested by diarrhea and dysentery [62]. However, it was reported that Saponin reduces body cholesterol by preventing its re-absorption and suppresses rumen protozoan by reacting with cholesterol in the protozoan cell membrane thereby causing it to lyse. It also binds to both exogenous and endogenous proteins including enzymes of the digestive tract, thereby affecting the utilization of protein [63, 64, 65].

The functional properties of the selected leafy vegetables are shown in Table 4 indicating the water absorption capacity (WAC), fat absorption capacity (FAC), emulsion capacity, emulsion stability, foaming capacity, least gelation concentration and foaming stability after 30 minutes. The WAC of the leafy vegetables ranges from 181.5 to 195.0 in *Ocimum gratissimum* to *Telfairia occidentalis* respectively. The FAC varied from 30.4 in *Vernonia amygdalina* to 38.2 in *Telfairia occidentalis*. The fat emulsion capacity and emulsion stability of the three leafy vegetables were in close range of 32.7 to 38.2 for emulsion capacity and 42.8 to 45.4 for emulsion stability. Also, foaming capacity, foaming stability and least gelation concentration were similar for the three vegetables (20.9 to 25.0 %, 4.4 to 4.6% and 7.8 to 8.3 % respectively). The functional properties (Table 4) shows that water absorption capacity is high and higher than the value reported for soybean (130%) [66], 134% reported by [67] for African yam bean but in close agreement with the work of Fasuyi [36], Ige [68] for melon seeds. WAC is an important protein property in viscous food such as gravies, soups, dough and

bakery products [67]. With these WAC values, it suggests that their incorporation into food formulation especially low protein traditional food such as maize-gruel, cassava flour and yam flour to enhance their nutritive values. Fat absorption capacity of the leafy vegetables were lower than the values reported for soybean and sunflower, 84.4 and 20% respectively [66] but in close agreement with work of [36]. Also, fat emulsion capacity values were higher than the values reported for soybean flour and sunflower (11.7 and 18%) respectively [66]. Fat absorption capacity is a critical factor in flavour retention while emulsion capacity is important for the stabilization of fat in the production of foods such as sausages, soups and cakes [69]. The ability of protein to aid the formation and stabilization of emulsion is very important in application such as mayonnaise, milk comminuted meat and salad dressing [67].

CONCLUSION

The results suggest that the vegetables if consumed in sufficient amount could contribute greatly towards meeting human nutritional requirement for normal body growth and adequate protection against diseases arising from malnutrition. Hence, they are highly recommended for consumption. These species contained substantial quantities of minerals like iron, calcium, phosphorus, potassium, magnesium and zinc. These species also contain high levels of some antinutritional factors like oxalic acids, polyphenols and Saponin, which are required to be removed to improve the nutritional quality of these leafy vegetables. However, values obtained for these vegetables are lower than the established toxic level. Hence they can be consumed without any restriction. However, consumption in large amounts of vegetables with higher levels of antinutrient should be avoided.

Table1: Proximate composition of different leafy vegetable species (100g/dry weight)

	O. gratissimum	V. amygdalina	T. occidentalis
Moisture (%)	84.0±0.1	80.12±0.2	79.92±0.1
Crude protein (g)	22.7±0.2	21.02±0.4	20.61±0.2
Total carbohydrate (g)	62.2 ±0.1	68.22±0.1	55.86±0.3
Crude fibre (g)	10.7±0.2	22.44±0.2	18.28±0.2
Ether extract	4.35±0.3	4.24±0.2	5.6±0.4
Ash	6.8±0.2	7.22±0.3	10.44±0.2
Fatty acids	1.17	3.65	4.82
Energy(kJ/100g)	1604.25	1673.96	1507.19

^acalculated fatty acids (0.86 x crude fat).

^bcalculated metabolizable energy (kJ/100g) (Protein x 17 + Fat x 37 + carbohydrate x 17)

Table 2: Mineral and vitamin content of species of leafy vegetable (mg/100 g)

	O. gratissimum	V. amygdalina	T. occidentalis
Sodium(Na)	11.4	12.6	14.4
Potassium (K)	35.2	44.8	48.8
Calcium (Ca)	15.4	15.8	18.7
Phosphorus (P)	13.8	13.1	15.08
Iron (Fe)	0.4	7.6	8.4
Magnesium (Mg)	12.2	14.2	18.7
Zinc (Zn)	3.2	1.2	1.4
Nickel	0.1	0.5	0.7
Na/k	0.32	0.28	0.30
Ca/P	1.12	1.21	1.24
Riboflavin	0.3	3.24	4.12

Vitamin C (mg/100g)	75.04	178.5	162.5
Niacin	1.2	1.3	1.4
Carotenoids (mg/100g)	28.0	35.0	48.0

Figures are mean of three replicates

Table 3. Concentration of some vitamins and antinutritional factors in three leafy vegetables

	<i>O. gratissimum</i>	<i>V. amygdalina</i>	<i>T. occidentalis</i>
Oxalate (mg/100g)	6.52±0.2	5.25±0.2	15.21±0.2
Saponin (mg/100g)	0.57±0.2	1.44±0.2	0.76±0.2
Polyphenol	10.12±0.2	6.68±0.2	5.52±0.2
Tannin	0.59±0.2	25.21±0.2	20.22±0.2
Phytate	0.63±0.2	22.24 ±0.2	18.40±0.2

Table 4 Amino acid contents of different leafy vegetable species(g/100 g) protein

Amino acids	<i>O. gratissimum</i>	<i>V. amygdalina</i>	<i>T. occidentalis</i>	FAO/WHO Pattern
Alanine	8.8	8.6	6.5	----
Aspartic acid	11.2	9.3	6.2	----
Cysteine	1.1	1.5	1.2	----
Arginine*	7.5	8.2	5.2	---
Methionine*	0.5	1.6	2.5	3.52
Glutamic acid	10.4	14.5	11.2	----
Histidine*	1.4	9.2	7.4	----
Glycine	4.1	6.6	6.2	----
Isoleucine*	3.8	4.7	5.2	4.00
Leucine*	7.1	7.2	7.6	7.04
Lysine*	3.1	3.4	2.4	5.44
Tyrosine	2.4	2.3	5.8	-----
Phenylalanine*	5.1	6.2	5.0	6.08
Threonine*	3.2	4.8	3.7	4.00
Tryptophan	1.6	1.5	3.2	1.00
Serine	4.4	5.2	4.0	----
Valine*	5.1	7.6	6.3	4.96

Figures are mean of three replicates

*Essential amino acids

Table 5. Functional properties of different leafy vegetable species

Amino acids	<i>O. gratissimum</i>	<i>V. amygdalina</i>	<i>T. occidentalis</i>
Water absorption capacity (%)	181.5±1.8	184.0±2.0	195.0±1.4
Fat absorption capacity (%)	34.2±0.8	30.4±3.0	38.2±1.7
Foaming capacity (%)	20.9±1.8	22.2±2.1	25.0±3.1
Foaming stability	43.5±1.2	42.8 ±2.4	45.4±2.2
Emulsion stability	4.4±0.1	4.5±1.2	4.6±3.1
Emulsion capacity (%)	32.7±0.4	38.2±1.8	35.1±2.1
Least gelatin concentration (%)	7.8±1.3	8.3±0.6	8.2±0.8

REFERENCES

1. FAO, 1985. Proceedings of the world Food survey conference. World Health Organization Tech. series Rome, Italy.
2. WHF, 2005. World Hunger Facts, World Hunger Education Service. In FAO World Food Summit Progress Report September 2004. FAO publications, Rome
3. Okafor, JC (1983). Horticultural Promising indigenous wild plant species of the Nigerian Forest zone. *Acta Hort.* 123: 165 – 176
4. Kala, A and Prakash 2004. Nutrient composition and sensory profile of differently cooked green leafy vegetables. *Intl. J. Food properties* 7: 659-669
5. Gupta, K., G.K. Barat, D.S. Wagle and H.K.L. Chawala, 1989. Nutrient contents and antinutritional factors in conventional and non conventional leafy vegetables. *Food Chem.*, 31: 105-116.
6. Oke, OL (1966). Chemical studies on the more commonly used vegetables in Nigeria. *Afr. Sci. Ass.* 11: 42-48.
7. Ladan MJ, Bilbils LS, Lawal M (1996). Nutrient composition of some green leafy vegetable consumed in Sokoto. *Niger. J. Basic Appl. Sci.* 5: 39-44.
8. Barminas, JT., M. Charles and D. Emmanuel, 1998. Mineral composition of non-conventional leafy Vegetables. *Plant Foods for Human Nutr.*, 53: 29-36.
9. Ifon, ET. and O. Basir, 1979. Nutritive value of some Nigerian leafy vegetables Part 3. *Food Chem.*, 5: 231-235.
10. Oshodi, AA. (1992) Proximate composition, nutritionally valuable minerals and functional properties of *adenopus breviflorus* bent seed flour. *Food Chem.*, Vol.45, pp.79-83.
11. Ejoh, A.R., M.F. Tchouanguép and E. Fokou, 1996. Nutrient composition of the leaves and flowers of *Colocasia esculenta* and the fruits of *Solanum melongena*. *Plant Food for Human Nutr.*, 49: 107-112
12. Devadas, RP. and S. Saroja, 1980. Availability of Fe and β -carotene from *amaranthus* to children. In: Emmaus P (Ed), *Proceeding of the 2nd Amaranthus Conference*, Rodale Press Inc., pp: 15-21.
13. Marfo, EK, Oke OL; and Afolabi, A, 1988. Nutritional evaluation of pawpaw (*carica papaya*) and flamboyant (*Delonix regia*) seed oils, *Nut. Res. Int.* 37(2), 303-310.
14. Udosen, EO. and Ukpanah, UM. (1993). Toxicants and Phosphorus content of some Nigerian vegetables. *Plant Food for Human Nutr.*, Vol. 44, pp. 285-289.
15. Aletor, VA and Adeogun 1995. Nutrient and antinutrient components of some tropical leafy vegetables. *Food chemistry*, 53: 375-379
16. Lockett CT, Calvert CC, Grivetti, LE (2000). Energy and micronutrient Composition of dietary and Medicinal wild plants Consumed during drought: Study of Rural Fulani, Northeastern Nigeria. *Int. J. Food Sci. Nutr.* 51: 195-208.
17. ECHO, 2003. Leafy vegetables. http://www.echonet.org/seeds/os_seed_catalogue/leafy-vegetables.htm
18. Akindahunsi, AA, and Salawu SO (2005). Photochemical screening and nutrient-anti-nutrient composition of selected tropical green vegetables. *Afr. J. Biotech.* 4: 497-501.
19. Edeoga, HO, Omosun G, Uche, LC (2006). Chemical composition of *Hyptis suaveolens* and *Ocimum gratissimum* hybrids from Nigeria. *Afr. J. biotechnol.* 5(10): 892-895.

20. Hassan LG, Umar KJ (2006). Nutritional value of Balsam Apple (*Momordica balsamina* L.) leaves. Pak. J. Nutr. 5(6): 522-529
21. Ekop, AS (2007). Determination of chemical composition of *Gnetum africana* (AFANG) seeds. Pak. J. Nutr. 6(1): 40-43.
22. Liener, IE (1969). Toxic constituents of plant foodstuff. Academic Press, New York, London.
23. Nwokolo, EN, Bragg, DB (1977). Influence of phytic acid and crude fibre on the availability of minerals from four protein supplements in growing chicks. *Can. J. Anim. Sci.* 57: 475-477.
24. Lewis J, Fenwick, GR (1987). Glucosinolate content of Brassica vegetables: Analysis of twenty-four cultivars of calabres. *Food Chemistry* 25: 259-268
25. A.O.A.C., 1995. Official methods of analysis of the Association of Analytical Chemists, 16th Edition. Washington D.C.
26. Jenson, N.F., 1978. Wild Plant fruit production: A Research symposium on Wild Food Plants
28. Igile, GO., W. Olesezk, S. Burda and M. Jurzysa, 1995. Nutritional assessment of *Vernonia amygdalina* leaves in growing mice. *J. Agric. Food Chem.*, 93:2162-2166.
29. Fevrier, C. and G. Viroben, 1996. La valeur nutritionnelle des matières protéiques végétales. In: Gordon B. Protéines végétales 2e édition. Tec. et Doc. Lavoisier, Paris, pp: 568-602.
30. Craplet, J. and Craplet-Meunier, 1979. Dictionnaire des aliments et de la nutrition. Edition Le Hameau, pp: 493.
31. Tanya, AKN., CMF. Mbofung and OO. Keshinro, 1997. Soluble and insoluble fiber contents of some Cameroonian foodstuffs. *Plant Foods for Human Nutr.*, 51: 199-207.
32. Eun-Hee, K., M. Young-Sun and W. Soon-Ja, 1993. Dietary fibre contents in some vegetables and seaweed's. *Korean J. Nutr.*, 26: 196-201.
33. Olatidoye, OP and Ogunshua, OA (2009). Production, Evaluation and Consumer Acceptability of *Ocimum* SPP as dry food seasoning. *Journal of research in Agriculture* Vol.6 No.3 pp8-14
34. Jenkin, DJ., AL. Jenkin, T.M.S. Wolever, A.V. Rao and L.U. Thompson, 1986. Fibre and starchy foods: gut function and implication in disease. *Am. J. Gastroenterol.*, 81: 920-930.
35. Singhal, R.S. and P.R. Kulkarni, 1987. Composition of the seeds of some Amaranths species *J. Sci. Food Agric.*, 42: 211-220
36. Fasuyi, O.A, 2006. Nutritional potentials of some tropical vegetable leaf meals: chemical characterization and functional properties *African Journal of Biotechnology* Vol. 5 (1), pp. 49-53,
37. FAO, 1986. Food composition table for use in Africa. Rome: FAO, Italy
38. WHO/FAO, 1985. WHO FAO report: Energy and protein requirements. WHO technical report series No 724. Geneva
39. Kutu, JO. and H.O. Kutu, 1999. Proximate composition and mineral content of two edible species of *Cnidioscolus* (tree spinach) *Plant Foods for Human Nutr.*, 53: 275-283.
40. Olaofe, O. Adeyemi. FO and Adediran, GO 1994. Amino acid and Mineral Composition and functional properties of some oil seeds. *J. Agric. Food Chem*, Vol. 42, pp.878-884

41. Oshodi, A.A., 1992. Comparison of proteins, minerals and vitamin C content of some dried leafy vegetables. Pak. J. Sci. Ind. Res., 35: 267-269.
42. Adeyeye, E (2004). The chemical composition of liquid and solid endosperm of ripe coconut. Orient. J. Chem. 20: 471-478.
43. Aremu MO, Olaofe O, Akintayo TE 2006. A comparative study on them chemical and amino acid composition of some Nigerian under-utilized Legume flours. Pak. J. Nutr. 5(1): 34-38
44. Kuri, Y.E., I. Sundar, K. Rao, L. Kahuwi, G.P. Jones and D.E. Rivelt, 1991. Chemical Composition of monardica characteristics, L. fruits. J. Agri. Food Chem., 39: 1702-1703.
45. Pineda, O., Torum, B., Viteri, F. and Arroyave, G.1981. In Bodwell, C.E. (ed.). Protein Quality in Humans. Avi Publishing Co., Westport, Conn., pp.12-23. .
46. Singh, G., Asha Kawatra and S. Sehgal, 2001. Nutritional composition of selected green leafy vegetables, herbs and carrots. Plant Foods for Human Nutr., 56: 359-364.
47. Lintas C (1992). Nutritional aspects of fruits and Vegetable Consumption. Options Mediterraennes 19: 79-87.
48. Wallece, PA., EK. Marfo and WA. Plahar, 1998. Nutritional quality and antinutritional composition of 4 non conventional leafy vegetables. Food Chem., 61: 287-291.
49. Deosthale, YG., VS. Mohan and KV. Rao, 1970. Varietal deficiencies in protein, lysine and leucine content of gram sorghum. J. Agric. Food Chem., 18: 644-646
50. Pingle, U. and B.V. Ramasastri, 1978. Absorption of calcium from leafy vegetables rich in oxalates. Br. J. Nutr., 39: 119-125
51. Oke OL (1969). Chemical studies on the more commonly used vegetables in Nigeria. Afr. Sci. Ass. 11: 42-48.
52. Vityakon, P. and B.R. Standal, 1989. Oxalate in vegetable amaranth (*Amaranthus gangeticus*). Forms, contents and their possible implications for human health. J. Sci. Food Agric., 48: 469-474.
53. Kikunaga, S., K. Ishii and M. Takahashi, 1995. The bioavailability of magnesium in spinach and the effect of oxalic acid on magnesium utilization examined in diets of magnesium deficient rats. J. Nutr. Sci. Vitaminol., 41: 671-68
55. Yadav, S Kala and SI. Sehgal, 2003. Effect of domestic processing and cooking on selected antinutrient contents of some green leafy vegetables Plant Foods for Human Nutr., 58: 1-11
56. Ridout, C.L., R.K. Price, M.S. Dupont, M.L. Parker and G.R. Fenwick, 1991. Quinoa saponins-analysis and preliminary investigations into the effects of reduction by processing. J. Sci. Food Agric., 54: 165- 176.
57. Koziol, M.J., 1990. Afrosimetric estimation of threshold saponin concentration for bitterness in quinoa. J. Sci. Food Agric., 54: 211-220.
58. Ladeji O, Akin CU, Umaru HA (2004). Level of antinutritional factors in vegetables commonly eaten in Nigeria. Afr. J. Nat. Sci. 7: 71-73.
59. Badifu GI, Okeke EM (1992). Effect of bleaching on oxalate hydrocyanic acid and saponin of four Nigerian leafy vegetables. J. Agric. Sci. Tech. 1: 71-75.

60. Erdman JN (1979) Oily seed phytates nutritional implications. J. Am. Oil Chem. Soc. (JOCS) 56: 736-741
61. Makkar HP, Becker K (1998). Plant toxins and detoxification methods to improve feed quality of tropical seeds. J. Animal Sci. 12: 467-480
62. Awe IS, Sodipo OA (2001). Purification of saponins of root of *Blighia sapida*. Niger. J. Biochem. Mol. Biol. 16(3): 201-204.
63. Bagepallis S, Narasinga R, Tatinemi P (1993). Tannin contents of foods commonly consumed in India and its influence on ionisable iron. J. Sci. Food Agric. 33: 89-96.
64. Aleto VA (1993). Allelochemical in plant food and feeding stuffs. Nutritional, biochemical and physiopathological aspects in animal production. Vet. Hum Toxicol. 35: 57-67.
65. Sotelo AE, Contreras S, Flores S (1995). Nutritional value and content of antinutritional compounds and toxics in ten wild legumes of Yucatan peninsula. Plant Food Hum. Nutr. 47: 115-123.
66. Lin, R.U., E.S. Humbert and F.W. Sosulki, 1974. Certain functional properties of Sunflower meal products. J. Food Sci., 39: 368-370
67. Adeyeye E, Otokili MKO (1999). Proximate composition and some nutritionally valuable minerals of two varieties of *Capsicum annum* (Bell and cherry peppers). Discov. Innov. 11: 75-81
68. Ige, M.N, A.O. Ogunsua and O.L. Okon, 1984. Functional properties of the protein of some Nigeria oil seeds. Casophor seeds and three varieties of some Nigeria oil seeds. Food Chem., 32: 822-825
69. Kinsella JE (1976). Functional properties of protein foods. Crit Rev. Food Sci. Nutr. 1, 219-236.